

CKDMIP progress meeting 8 Sept 2020

1. What is required from participants?
2. Results so far
5. Questions for discussion

Robin Hogan and Marco Matricardi

ECMWF

r.j.hogan@ecmwf.int

Any objections to recording this meeting for the benefit of colleagues who could not attend? Recording will not be posted publicly on internet.

Agenda


1. What is being asked of CKDMIP participants? (Robin)
 - Discussion: what further help/software/data would be of use to participants?
2. Summary of results so far (Robin)
3. Overview of SOCRATES CKD tool (James Manners)
 - Discussion: what are the interesting differences between CKD approaches that we can evaluate using the CKDMIP dataset?
4. Towards a community tool chain for gas optics (Robert Pincus)
 - Discussion: what are the community needs, can existing codes be released open source?
5. Discussion of next steps (all)
 - What is a realistic timeline: from gathering contributions from all models to a CKDMIP results paper?
 - What else is needed, e.g. clouds, spectrally varying surface albedo, non-LTE...?
 - Next meeting(s)?

Objectives

- To use benchmark line-by-line calculations to evaluate the accuracy of existing CKD models. *Done for RRTMG, RRTMGP and RRTMGP-NN (LW)*
- To explore how accuracy varies with number of k-terms / g-points in individual CKD schemes for applications spanning short-range weather forecasting to climate modelling. *Done for ecCKD*
- To understand how different choices in way that CKD models are generated affects their accuracy for the same number of g-points.
- To provide freely available datasets and software to facilitate the development of new gas-optics models, with the ultimate aim of producing a community tool to allow users to generate their own gas-optics models targeted at specific applications.

100 evaluation profiles of T , q and $[O_3]$ in two evaluation datasets

- *Evaluation-1*: 50 profiles for which the LBL spectra and broadband fluxes are available to participants
- *Evaluation-2*: 50 profiles for which the spectra and fluxes are withheld for independent evaluation




Trace gas variations: 34 scenarios (18 in shortwave)



[justification](#)

Scenario	Comment	CO ₂ ppmv	CH ₄ ppbv	N ₂ O ppbv	CFC-11 eq. pptv	CFC-12 pptv
1	Glacial maximum	180	350	190	32	0
2	Preindustrial	280	700	270	32	0
3	Present-day (2020)	415	1921	332	861	495
4	Future (2110)	1120	3500	405	2000	200
5–9	CO ₂ forcing	180, 280, 560, 1120, 2240	1921	332	861	495
10–14	CH ₄ forcing	415	350, 700, 1200, 2600, 3500	332	861	495
15–18	N ₂ O forcing	415	1921	190, 270, 405, 540	861	495
*19–20	CFC-11 forcing	415	1921	332	0, 2000	495
*21–22	CFC-12 forcing	415	1921	332	861	0, 550
*23–24	CO ₂ /CH ₄ overlap	180, 2240	350	332	861	495
*25–26		180, 2240	3500	332	861	495
*27–28	CO ₂ /N ₂ O overlap	180, 2240	1921	190	861	495
*29–30		180, 2240	1921	540	861	495
*31–32	CH ₄ /N ₂ O overlap	415	350, 3500	190	861	495
*33–34		415	350, 3500	540	861	495

- One concentration file per scenario, containing 50 profiles
- 2x50x34 = 3400 profiles in 2x34 = 68 files, e.g. **ckdmip_evaluation1_concentrations_present.nc**



Participants requested to use 5 "wide" and 13 "narrow" bands in SW and LW, but single band (FSCK) also permitted



Line-by-line absorptions calculated by Marco Matricardi

SW workflow for one scenario, one CKD model

ckdmip software package
mean-ssi_nrl2.nc
Solar spectral irradiance

FTP site
ckdmip_evaluation1_concentrations_present.nc
50 profiles of T , p and mole fractions of 9 gases in layers and at layer interfaces

Participants' tasks

Your shortwave CKD model for a particular application, band structure and total number of g-points

One file per CKD model
ecckd-0.6_sw_climate_wide-38_spectral_definition.nc
• `gpoint_fraction(wavenumber_interval, g_point)`

One file per CKD model/scenario
ecckd-0.6_evaluation1_sw_climate_wide-38_optical-depth_present.nc
• `optical_depth(column, layer, g_point)` *or total optical depth*
• `rayleigh_optical_depth(column, layer, g_point)` *& single scat. albedo*
• `incoming_sw(column, g_point)`

FTP site
ckdmip_evaluation1_sw_fluxes_present.nc
LBL fluxes

ckdmip_sw ckd mip software package

ecckd-0.6_evaluation1_sw_climate_wide-38_fluxes_present.nc

ckdmip software package
Matlab scripts

Evaluation plots on CKDMIP web site

LW workflow for one scenario, one CKD model

Participants' tasks

Your **longwave** CKD model for a particular application, band structure and total number of g-points

One file per CKD model

ecckd-0.6_lw_climate_wide-42_spectral_definition.nc
• gpoint_fraction(column, layer, g_point)

ecckd-0.6_evaluation1_lw_climate_wide-42_optical-depth_present.nc
• optical_depth(column, layer, g_point)
• planck_hl(column, half_level, g_point)

One file per CKD model/scenario

ckdmip_lw

ckdmip software package

FTP site

ckdmip_evaluation1_lw_fluxes-4angle_present.nc
LBL fluxes

ecckd-0.6_evaluation1_lw_climate_wide-42_fluxes-4angle_present.nc

ckdmip software package

Matlab scripts

Plots on CKDMIP web site

FTP site

ckdmip_evaluation1_concentrations_present.nc
50 profiles of T , p and mole fractions of 9 gases in layers and at layer interfaces

How many CKD models should I submit?

- Can be as few as one, e.g. RRTMG, RRTMGP, Fu-Liou...
- If it affects the number of g points you need, you can vary the [application](#):
 - Climate (all scenarios, 0.02-1100 hPa), Global NWP (present-day scenario only, 0.02-1100 hPa), Limited-area NWP (present-day scenario, 4-1100 hPa)
 - Evaluation statistics will be limited to applicable scenarios and pressure ranges
- You can vary the [band structure](#):
 - CKDMIP proposes a set of *narrow* and *wide* bands, but other band structures accepted
- You can vary the number of g points (k terms)
 - Explore the trade-off between accuracy and efficiency
- Could be $3 \times 2 \times 6 = 36$ different CKD models! Up to you how many you submit depending on interest (e.g. only climate) and your time

What do I do with the LBL datasets?

- If you can only use your own LBL model to generate CKD models, you can ignore them
- However, intention of CKDMIP is to compare CKD models generating assuming the *same* spectroscopy, so we can eliminate spectroscopy differences as a cause of apparent errors
- Therefore, the FTP site contains absorption spectra for three of the four CKDMIP datasets (*Evaluation-2* held back for independent evaluation) using LBLRTM 12.8, which you can use for training (ecCKD uses all three of these datasets, for example)


Name	Purpose	Layers	T profiles	Description
<i>Evaluation-1</i>	Training & evaluation	54	50	Realistic profiles selected from NWP-SAF dataset
<i>Evaluation-2</i>	Independent evaluation	54	50	Further profiles selected from NWP-SAF dataset
<i>MMM</i>	Training	52	3	Median, min. and max. of NWP-SAF T , q and O_3 profiles
<i>Idealized</i>	Generating look-up tables	53	11	Idealized profiles regularly spaced in T , $\log p$ and $\log q$

- The Technical Guide explains how the CKDMIP software can be used to perform some manipulation of these datasets, including radiative transfer calculations
- See [contents of FTP site](#)


What does the “spectral definition” file contain?

- Normally we have N g-points in a band but don't care *where* in the band
- For wide bands this becomes important when representing clouds: for more accuracy the cloud optical properties could be parameterized per g point not per band
- The spectral definition file contains `gpoint_fraction`, which sums to 1 along the wavenumber axis
- If your model uses a different $g(w_n)$ relationship at different pressures, I suggest you use a mid-tropospheric reference pressure of 500 hPa
- [Jump to end](#)


Longwave ecCKD wide-43 model









Shortwave ecCKD wide-38 model



Use realistic vertical profiles for the five well-mixed greenhouse gases



CKDMIP Evaluation-1 dataset: Well-mixed gases



- Consider five well-mixed greenhouse gases
- Further 38 implicitly considered via “equivalent CFC11” (Meinshausen et al. 2017)
- Evaluate concentrations from Glacial Maximum to the worst of the CMIP6 future scenarios
- Participants produce CKD models for both climate and present-day NWP
- Per-molecule absorption of these gases is independent of concentration, so only store absorption spectra at one concentration and scale
- Also N₂ and O₂ absorption

Three “applications”

Application	Lowest pressure	GHG concentrations
Limited-area NWP	4 hPa	Present-day (2020)
Global NWP	0.02 hPa	Present-day (2020)
Climate	0.02 hPa	Variable

- Most contributors are interested primarily in the “climate” application: your model will be tested over all 34 climate scenarios at pressures down to 0.02 hPa
- If you think you can produce a faster CKD model (fewer g points) for present-day NWP applications, you can submit results for separate CKD models that will be evaluated only using the “present” greenhouse-gas scenario, and optionally also only pressures down to 4 hPa for limited-area NWP

Two proposed band structures: 13 narrow bands, 5 wide bands

Longwave

<i>Narrow bands</i>			<i>Wide bands</i>	
#	Spectral interval (cm ⁻¹)	RRTMG <i>k</i> terms	#	Label
1	0–350	8	1	Far infrared
2	350–500	14		
3	500–630	16	2	Main CO ₂ band
4	630–700	14		
5	700–820	16		
6	820–980	8	3	Infrared window
7	980–1080	12		
8	1080–1180	8		
9	1180–1390	12	4	Mid-infrared A
10	1390–1480	6		
11	1480–1800	8		
12	1800–2080	8	5	Mid-infrared B
13	2080–3260	10		

Shortwave


<i>Narrow bands</i>			<i>Wide bands</i>	
#	Spectral interval (cm ⁻¹)	RRTMG <i>k</i> terms	#	Label
1	250–2600	12	1	Mid-infrared
2	2600–3250	6		
3	3250–4000	12		
4	4000–4650	8	2	Shortwave infrared
5	4650–5150	8		
6	5150–6150	10		
7	6150–8050	12		
8	8050–12850	10		
9	12850–16000	8	3	Near infrared
10	16000–22650	6	4	Visible window
11	22650–29000	6		
12	29000–38000	8	5	Ultraviolet
13	38000–50000	6		

FTP site directory contents

- **concentrations/** 34 scenarios for each of the Evaluation-1 and Evaluation-2 datasets
 - **lw_spectra/, sw_spectra/** LBL gas absorption spectra (big!)
 - **evaluation1/** Spectra for *Evaluation-1* dataset (*Evaluation-2* kept back for independent evaluation)
 - **idealized/** Spectra for regularly spaced temperature pressure and humidity – can use for training
 - **mmm/** Spectra for maximum, minimum and median T, q and [O3] – can use for training
 - **lw_fluxes/, sw_fluxes/:** LBL fluxes per narrow band for each scenario & profile – for evaluation
 - **evaluation1/**
 - **results/**
 - **CKD-TOOL/lw_optical-depth/, CKD-TOOL/sw_optical-depth/** Files submitted by participants
 - **CKD-TOOL/lw_fluxes/, CKD-TOOL/sw_fluxes/** Fluxes calculated by from submitted optical depths
- Not essential!
Available for participants if they wish

Dataset volumes

- Absorption spectra use 7.1M points in longwave, 3.1M in shortwave
- Stored in NetCDF4/HDF5 format with compression
- *Evaluation-1 dataset:*
 - 50 profiles: 265 GB in LW, 136 GB in SW
- *Evaluation-2 dataset:*
 - For independent evaluation (not released)
 - Same size as Evaluation-1
- *MMM dataset:*
 - Median, min and max profiles
 - 3 profiles: 30 GB in LW, 15 GB in SW
- *Idealized dataset:*
 - For creating CKD look-up tables
 - 6 profiles: 139 GB in LW, 74 GB in SW



End of part 1

Results









Models and tools participating in CKDMIP

CKD model	Contact	LW status	SW status
RRTMG (ecRad)	Robin Hogan	v1.2.0	v1.2.0
RRTMGP (RTE)	Robert Pincus	v181204	v181204
RRTMGP-NN	Peter Ukkonen	v1.0	
MODTRAN	Alexander Berk		
Fu-Liou	Lei Lin		

CKD tool	Contact	LW status	SW status
SOCRATES	James Manners		
ecCKD	Robin Hogan	v0.5	v0.6
ARTDECO-PyKdis	Mathieu Compiegne		
INSA-CNRS/LOA/HYGEOS	Frederic Andre, Mathieu Compiegne		
PSLACKD	Seiji Kato, Fred Rose		
CMA scheme	Hua Zhang		
mstrnX	Miho Sekiguchi		
KBIN	Nils Madenach, Juergen Fischer, Rene Preusker		

Longwave RRTMG

- 16 bands
- 140 g points



Scenario: Present-day (2020)

CKD model: ecRad-RRTMG

Bias TOA upwelling: -0.35 W m^{-2}

Bias surface downwelling: -0.40 W m^{-2}

RMSE TOA upwelling: 0.59 W m^{-2}









RMSE surface downwelling: 0.73 W m^{-2}

RMSE heating rate (0.02-4 hPa): 0.194 K d^{-1}

RMSE heating rate (4-1100 hPa): 0.106 K d^{-1}

Longwave RRTMGP

- 16 bands
- 256 g points



Scenario: Present-day (2020)

CKD model: RTE-RRTMGP

Bias TOA upwelling: -0.43 W m^{-2}


Bias surface downwelling: 0.20 W m^{-2}

RMSE TOA upwelling: 0.63 W m^{-2}



RMSE surface downwelling: 0.63 W m^{-2}


RMSE heating rate (0.02-4 hPa): 0.127 K d^{-1}

RMSE heating rate (4-1100 hPa): 0.099 K d^{-1}





Longwave RRTMG

















Longwave RRTMGP



Longwave RRTMG











Longwave RRTMGDO



Shortwave RRTMG

- 14 bands
- 112 g points



Scenario: Present-day (2020)

CKD model: ecRad-RRTMG (fix SSI)

Bias TOA upwelling: 0.68 W m^{-2}

Bias surface downwelling: 1.75 W m^{-2}

RMSE TOA upwelling: 0.76 W m^{-2}









RMSE surface downwelling: 1.96 W m^{-2}

RMSE heating rate (0.02-4 hPa): 0.660 K d^{-1}

RMSE heating rate (4-1100 hPa): 0.136 K d^{-1}

Shortwave RRTMGP

- 14 bands
- 224 g points



Scenario: Present-day (2020)

CKD model: RTE-RRTMGP

Bias TOA upwelling: -0.06 W m^{-2}

Bias surface downwelling: -0.05 W m^{-2}


RMSE TOA upwelling: 0.14 W m^{-2}

RMSE surface downwelling: 0.73 W m^{-2}


RMSE heating rate (0.02-4 hPa): 0.256 K d^{-1}

RMSE heating rate (4-1100 hPa): 0.056 K d^{-1}


Shortwave RRTMG




Shortwave RRTMGP



Shortwave RRTMG




Shortwave RRTMGP




ecCKD: error versus computational cost (number of k terms) for all scenarios

- Unsurprisingly, error decreases with number of k terms, but can flatten off
- **Full-spectrum correlated- k (FSCK) method** works well in longwave, but not yet in shortwave


Longwave:




Shortwave:



[See detailed plots](#)







mwf.int/display/C/



More efficient to train for NWP applications?

- Rather than all **climate** scenarios, **global-nwp** trains for present-day only, and low-top **limited-area-nwp** for pressures larger than 4 hPa (rather than 0.02 hPa)

Longwave:







Shortwave:

Low-top model doesn't waste k terms for accurate heating in upper atmosphere, so gets same accuracy in shortwave fluxes with fewer k terms

Not much difference for global-nwp, or longwave

wrf.int/display/C



End of part 2

Discussion of next steps

- GMD protocol paper has just received second review, will be revised in the coming few weeks, and one of the reviewers raised a question of the scope of CKDMIP: [should it include the effect of uncertainties in spectroscopy?](#)
- What is a realistic timeline for gathering contributions from all models?
- What is a realistic timeline to a CKDMIP results paper?
- Making things more complicated (may be possible without more data from participants)
 - Clouds & aerosols
 - Spectrally varying surface albedo and emissivity
 - Non-LTE?
- Other science questions that could be answered?
- Next meeting: date and purpose?


Should CKDMIP tackle the impact of spectroscopic uncertainties?


Reviewer comment: *The design of the CKDMIP, particularly the first and second evaluation data sets (table 1 and 2) containing **randomly selected realistic thermodynamics profiles, is going to make it much harder to search for systematic errors associated with the treatment of water vapor than is necessary** or feasible. Since water vapor is by far the dominant radiatively active species in both the shortwave and the longwave, and since the literature cited in the first major comment above shows that there is still **significant spread in the accuracy of the parameterization of near-IR H₂O absorption** across the CMIP ensembles, this is my principal concern regarding the design of CKDMIP. It's really important to be able to look at the change in the k distributions and resulting fluxes and heating rates when water vapor alone is perturbed. There is a simple fix for this, fortunately, **if CKDMIP were to also ask for the exact same set of data from each contributing group for the idealized profiles** as for the Evaluation-1 and, ultimately, Evaluation-2 datasets.*

- Should CKDMIP be limited to algorithmic problem of formulating a CKD model *given a spectroscopic database*? Previous intercomparisons have always mixed algorithmic and spectroscopic differences
- Should uncertainties in spectroscopy (especially in water vapour) be introduced into CKDMIP?
- Will they be present in CKDMIP inadvertently because some participants are not using the same spectroscopy, and what can we do about it?

Longwave ecCKD









- climate-fsck-27





Shortwave ecCKD

- climate-wide-38



Scenario: Present-day (2020)
CKD model: ecCKD-wide-38

Bias TOA upwelling: 0.52 W m^{-2}

Bias surface downwelling: -0.08 W m^{-2}

RMSE TOA upwelling: 0.64 W m^{-2}

RMSE surface downwelling: 0.73 W m^{-2}

RMSE heating rate (0.02-4 hPa): 0.202 K d^{-1}

RMSE heating rate (4-1100 hPa): 0.059 K d^{-1}

