

MARS

Introduction and basic concepts

Computer User Training Course 2014

Carsten Maaß

User Support

Contents

- **Introduction**
- **Meteorological content**
- **MARS language**
- **MARS architecture**
- **Retrieving data**
- **Practicals**

Introduction

Meteorological Archival and *Retrieval System*

- Meteorological data (**GRIB**: fields, **BUFR**, **ODB**: observations)
- Large amount of data (size of archive & number of fields)
- Operational & Research environment
- Batch & interactive modes
- Large number of users with different requirements:
 - large datasets rarely \leftrightarrow few fields very often
- Heterogeneous environment

Introduction – MARS components

- **Client/Server architecture**
- **Clients: workstations, supercomputers**
- **Servers: supercomputers, dedicated servers**
- **Several databases**
- **Tape library**

Introduction – Some figures

- **43 PetaBytes of data in ~ 10 million files, for more than 140 billion ($1.4 \cdot 10^9$) meteorological fields**
- **~ 280 Gigabytes of metadata**
- **$200 \cdot 10^6$ fields added daily (65 Terabytes)**
- **650 active users/day executing 1.5 million requests/day**
- **~ 100 Terabytes retrieved daily**
- **Analysis, forecast and observations since 1957 (ERA-40)**
- **Operational forecast since 1985**

Terminology – Forecast lead times

- **Medium-range**
 - the high-resolution and the ensemble forecasts of weather, at the space and time-scales represented by the relevant model, up to 10 and 15 days ahead, respectively, and the associated uncertainty
- **Extended-range (monthly)**
 - ensembles of individual forecasts and post-processed products of average conditions (e.g. weekly averages) up to 1 month ahead, and the associated uncertainty
- **Long-range (SEAS)**
 - ensembles of individual forecasts and post-processed products of average conditions (e.g. monthly averages) up to 13 months ahead, and the associated uncertainty

Terminology – ... more

- **Re-forecasts**

forecasts run for past decades necessary to estimate the model climate and the level of skill and to generate some of the operational products

- **IFS**

`Integrated Forecasting System`, *the system used at ECMWF*

<http://www.ecmwf.int/about/forecasts.html>

Meteorological content – Operational Analyses

- **4DVAR (T1279 / 16km, T255 inner loops, input to HRES)**
 - At synoptic hours 00, 06, 12 and 18 UTC
 - Surface
 - Model levels (137)
 - Pressure levels (25)
 - Isentropic levels (15 PT, 1 PV)
- **EDA (T399 / 50 km, T159 inner loops, input to ENS)**
 - At synoptic hours 00, 06, 12 and 18 UTC
 - 26 members
 - Surface
 - Model levels (137)
 - Pressure levels (25)
 - Isentropic levels (16 PT, 1 PV)

Meteorological content – HRES

- Atmospheric Forecast (10 day forecast based on 00/12 UTC Analysis) at T1279L137 (16 km)
 - Surface
 - Model levels (137)
 - Pressure levels (25)
 - Isentropic levels (16 PT, 1 PV)
 - 1 hourly steps from 0 to 90, 3 hourly from 93 to 144 and 6 hourly from 150 to 240 hours

Meteorological content – ENS

- Medium-range forecasts to 15 days, 91 Levels
- 26 member Ensemble of Data Assimilations (EDA, stream elda)
- 1 control forecast (as HRES but with lower resolution)
- 50 different forecasts with perturbed initial conditions
- Truncation at day 10 from T639 (~32 km) to T319 (~64 km)
- Two additional calibration/validation runs were added
- Leg 3: 00 UTC FC extended Mondays & Thursdays to day 32

	# FC	Leg 1 day 0-10	Leg 2 day 11-15	Leg 3 day 16-32
ENS-CF	1	T639		T319
ENS-PF	50	T639		T319
CV-T639	1	T639		
CV-T319	1	T319		

www.ecmwf.int/products/changes/vareps-monthly/

Meteorological content – ENS products

- **Control forecast**
- **Calibration/Validation forecasts**
- **50 perturbed forecasts / 26 EDA members**
- **Initial condition perturbations**
- **Ensemble mean and standard deviation**
- **Extreme forecast index**
- **Event probabilities**
- **Cluster mean, cluster representatives and standard deviation**
- **Trajectories (of tropical cyclones)**

Meteorological content – Ocean-Wave component

	Forecast/ Analysis	Domain	Number of members	Horizontal resolution	Number of directions	Number of frequencies
LAM WAM (LAW)	Analysis + forecast 0–5 days	Limited: 5° N–90° N, 98° W–54° E	1	11 km	36	36
WAM HRES	Analysis and forecast 0–10 days	Global	1	28 km	36	36
WAM ENS	Forecast 0–10 days	Global	51	55 km	24	30
WAM ENS	Forecast 10–32 days	Global	51	55 km	12	25
WAM SEAS	Forecast 0–13months	Global	51	111 km	12	25

Meteorological content – BC

Boundary condition forecast (Short cut-off forecast T1279L137 at 06/18)

- **Analysis (4DVAR)**
- **Forecast (to 90 hours) in hourly steps**
- **00/12 UTC AN/FC is taken from HRES**
- **Full fields are available**

Valid data only available for participating MS.

Meteorological content – Seasonal System 4

SEAS – atmosphere-ocean coupled model (51 members)

- Global forecasts from 00 UTC to 7 months: (once a month)
 - atmosphere: ~75 km resolution, 91 levels (T255 L91)
 - ocean: NEMO – ORCA1 grid (~ $1^\circ \times 1^\circ$ with equatorial refinement), 42 levels
- In February, May, August and November, 15 of the 51 members are extended to 13 months
- Re-forecasts: 15 members (0-13m) covering 30 years (1981-2010)
- Part of the EUROSIP system, with UK Met Office, Météo France and NCEP
- Availability of products: 12:00 on the 8th of each month

See www.ecmwf.products/changes/system4/

Meteorological content – Monthly Means

Averaged over each calendar month

- **Atmosphere / Wave**

- Analysis
- Forecast

Meteorological content – Special datasets (1/2)

- **Special Projects**

- ECMWF Re-Analyses (**ERA-Interim**, ERA-40, ERA-15)
- DEMETER: Multimodel Ensemble for seasonal to Interannual prediction
- Data targeting system
- ENSEMBLES
- EURO4M
- MACC
- PROVOST
- ECSN-Hyretics

Meteorological content – Special datasets (2/2)

- **IFS Research experiments**

- ECMWF
- Member States

- **Member States' Projects**


- COSMO-LEPS
- Aladin-LEAF

Meteorological content – ERA-Interim

- 35 years (1/1979 – 12/2013) of validated ERA-Interim analysis products are available
- Continued in near real-time (with ~2 months delay)
- Uses IFS Cycle 31r2, and 12h-4DVar
- Resolution:
 - Horizontal: T255, N128 (~0.7°)
 - Vertical: 60 ML, 37 PL, 16 PT, PV=±2
- Analyses at 00/06/12/18, Forecasts at 00/12 to 240 h
- Subset of products now also publicly available on the ECMWF Data Server at **full** resolution

ERA-Interim skill

- Uses IFS Cycle 31r2, T255L60 and 12h-4DVar
 - Improved model physics
 - New humidity analysis
 - Better formulation of background error constraint
 - Variational bias correction of satellite radiance data
 - Reprocessed Meteosat winds and clear sky radiances
 - ...



Meteorological content – TIGGE

- **THORPEX Interactive Grand Global Ensemble**
- **Global ensemble forecasts to around 14 days generated routinely at different centres around the world**

ECMWF, JMA (Japan), Met Office (UK), CMA (China), NCEP (USA),
MSC (Canada), Météo-France (France), BOM (Australia), CPTEC (Brazil),
KMA (Korea)

- Data archived in GRIB 2
- TIGGE-LAM data since 1/1/2013

<http://tigge.ecmwf.int>

Meteorological content – Observations & Feedback

- **Observations**

- Surface data
- Vertical soundings
- Upper-air data
- Satellite

- **Feedback**

- **Analysis Input**

- **Analysis Feedback (superseded by ODB feedback)**

Meteorological content – Data formats

WMO formats

- Fields in GRIB (**GRid In Binary**), ECMWF local extensions
 - Spherical Harmonics (upper-air fields, T1279)
 - Gaussian Grid (surface fields, N640)
 - Latitude/Longitude (wave and ocean products)
- Observations in BUFR (**Binary Universal Form Representation**)
 - Instrument specific

ECMWF/IFS format

- ODB (**Observational Data Base**)
 - Observation feedback

MARS – ODB

- In the IFS observations are handled by ODB (Observational Data Base)
- ODB is a
 - Hierarchical in-core database with a data definition and query language: ODB/SQL
 - A data format
 - ...
- ODB Observation Feedback (ofb) data is archived in MARS
 - Improve the representation of feedback data in MARS meta data
 - Introduce SQL capabilities to request feedback data
- To improve the handling of observations, ODB will be further integrated into ECMWF systems

MARS – future development

- **Content**

- **Multi-layer surface fields**
- **ERA-CLIM (<http://www.era-clim.eu>)**
- **JRA-55**

- **Architecture**

- **New interpolation package (MIR)**
- **Align with new Product Generation**

MARS language

Mechanism to *name* archived fields

Request syntax:

```
verb,  
      keyword1      = value1,  
      ...           = value2,  
      keywordN      = valueN
```

- **verb:** action to be taken (e.g. retrieve, list, read)
- **keyword:** a known MARS variable, e.g. type or date
- **value:** value assigned to the keyword, e.g. Analysis or temperature

MARS language

- **verb** and **keyword=value** separated by commas, but last one
- Spaces and tab characters are ignored
- *, ! and # comment until end-of-line
- Directives are not case sensitive
- Values: predefined names, numeric values or strings (filenames)
- Abbreviations: enough letters to uniquely identify keyword or value
- Acronyms: usually initial letters of names
- / is used as list separator → specify pathnames in quotes

MARS language – Retrieve request

retrieve ,		action
class	= od ,	identification
stream	= oper ,	
expver	= 1 ,	
date	= -3 ,	date & time related
time	= 12 ,	
type	= analysis ,	data related
levtype	= model levels ,	
levelist	= 1/to/137 ,	
param	= temperature ,	
grid	= 2.5/2.5 ,	post-processing
target	= “analysis”	storage

MARS language – Identification of archive

- class** ECMWF classification (od, rd, e4, ...)
- stream** originating forecasting system (oper, wave, enfo, seas, ...)
- expver** version of the experiment (01 operational, 11, aaaa)
- domain** area covered by the data (Global, Mediterranean, ...)
- origin** originating centre of the data (kwbc, egrr, ...)
- system** seasonal forecast operational system (1, 2, 3)
- method** to specify how the seasonal forecast is produced, e.g. in System 2, method=0 for runs without ocean assimilation (0, 1, ..., 3)

MARS language - Date & time

<input type="checkbox"/> time	base time or observation time (00, 06, 09:30, ...)
<input type="checkbox"/> date	base date of the model (-1, 20010225, ...)
<input type="checkbox"/> step	forecast time-step from base time (12, 24, 240, ...)
reference	reference forecast time step for EPS tube (96,...)
refdate	date of real-time forecast associated to re-forecast/hindcast (stream=mnfh)
hdate	base date of a re-forecast/hindcast (stream=enfh)
range	observations: period in minutes from base time (360,...) ocean fields: extension of the time series/average
fcmonth	month from seasonal forecast base date (1, 6, ...)
fcperiod	period, in days, for an averaged field (26-32)

MARS language – Fields

<input type="checkbox"/> type	type of field (an, fc, ...)
<input type="checkbox"/> levtype	type of level (pl, ml, sfc, pt, pv)
<input type="checkbox"/> levelist	levels for the specified levtype (off if levtype=sfc)
<input type="checkbox"/> param	meteorological parameter (t, temperature, 130, 30.128)
number	ensemble member (1, 2, ...)
channel	brightness temperature frequency band
diagnostic, iteration	sensitivity forecast products
frequency, direction	2-d wave spectra products
product, section, latitude, longitude	ocean products

MARS language – Observations & images

type

type of observations or images (ob, fb, ai, af, im)

obstype

observation subtype (s, air) or image channel

ident

WMO observation station number or satellite identifier

duplicates

whether duplicated observations are to be kept or not

block

WMO block number for observation

MARS language – ODB

reportype classification to index ODB data (16020)

filter SQL filter query ("select lat,lon,obsvalue where varno=39")

MARS language – Storage

- target** UNIX pathname where retrieved data is stored
- source** UNIX pathname from where to read data
- fieldset** temporary storage; can be considered a MARS variable

Unix pathnames (using /) have to be enclosed in quotes, e.g.

```
target = “/scratch/ms/gb/uid/analysis”
```

MARS language - Post-processing (1/2)

grid

output grid mesh

- Latitude/longitude increments in degrees (2.5/2.5)
- Number of latitude lines from Pole to Equator (160)

gaussian

type of Gaussian grid (regular, reduced)

area

desired sub-area in degrees (north/west/south/east)

frame

number of grid points from sub area inwards (5)

resol

triangular truncation (319, auto, av)

MARS language - Post-processing (2/2)

rotation **lat/lon of South Pole**

accuracy **number of bits per data value in GRIB (16)**

style **specify post-processing style (dissemination)**

MARS language – Execution control

expect **number of expected fields (1000, any, ...)**

database **where to look for the data**

use **hint about frequency of use (infrequent)**

MARS language – Values

- Single value, predefined names, numbers, mnemonics

`param = temperature`

- List of values, separated by `/`

`step = 12/24/48`

- Range of values, using keywords: `to`, `/` and `by`

`date = 20020101/to/20020131`

`step = 24/to/240/by/24`

MARS language – Values

- Expected number of fields is computed by multiplying number of values after expansion of ranges

`date = 20020101/to/20020131` 31 fields

- Certain keywords accept **all** as valid value

`levelist = all`

- Most keywords accept **off** as valid value

`levtype = surface,`

`levelist = off`

- Not all possible combinations **keyword = value** name an archived field

Request examples – Interim Re-Analysis

Retrieval of snow depth from the ERA-Interim archive for December 2007, for all analysis base times. It retrieves 124 fields.

```
retrieve,  
    class      = ei,  
    stream     = oper,  
    expver     = 1,  
    date       = 20071201/to/20071231,  
    time       = 00/06/12/18,  
    type       = an,  
    levtype    = sfc,  
    param      = sd,  
    target     = "era-int.200712.sd"
```

Request examples - Ensemble forecast

Retrieval of surface temperature and 10-m wind components (U and V), 20 first members of the EPS for 2nd Jan 2001 for time steps 12, 36 and 60. It retrieves 180 fields.

```
retrieve,  
        class      = od,  
        stream     = enfo,  
        expver    = 1,  
        date       = 20010102,  
        time       = 12,  
        step       = 12/36/60,  
        type       = pf,  
        levtype   = sfc,  
        param     = st/10u/10v,  
        number    = 1/to/20,  
        target    = "perturbed.sfc"
```

Request examples – Operational analysis

Retrieval of sea surface temperature for first 10 days of May 2002, all synoptic times. It retrieves 40 fields.

```
retrieve,  
        class      = od,  
        stream     = oper,  
        expver    = 1,  
        date       = 20020501/to/20020510,  
        time       = 00/06/12/18,  
        type       = an,  
        levtype   = sfc,  
        param     = sea surface temperature,  
        target    = "sst"
```

Retrieving data – Calling MARS

- **directives from input stream**

```
mars <<EOF
retrieve,
type = an,
date = -1,
target = "$SCRATCH/my_an"
EOF
```

- **directives from file**

```
cat > my_request <<EOF
retrieve,
type = an,
date = -1,
target = "$SCRATCH/my_an"
EOF
mars my_request
```

MARS Practicals

Point your browser to

www.ecmwf.int/publications/manuals/mars/practice/

or navigate to

Publications > Manuals > MARS > Practicals and examples

and follow the instructions

Retrieving data – Hints

- Default values: minimize their use
- No semantic check (only syntax is checked)
- MARS messages
 - INFO request execution and report
 - WARNING unusual aspect of execution
 - ERROR system or data errors
 - FATAL terminates execution

Web-MARS – <http://apps.ecmwf.int/services/mars/catalogue/>

Web interface to archive catalogue

- Content browsing of **every field in the archive**
- Real-time (dynamic access to metadata)
- Create MARS requests (without checking availability)
- Check availability of data
- Retrieval in GRIB and NetCDF for few fields
- URL based in MARS requests (can be edited & bookmarked)
- More up to date than static content documentation

Additional features in “old” Web-MARS (1/2)

<http://www.ecmwf.int/services/archive/>

● Data Finder

- Allows to have different views of the archive
 - By period of time
 - By meteorological parameter
 - By data source (IFS configuration)
- Narrow the search for data
- Brings you directly into the catalogue

● Changes in the archive

- Addition or discontinuation of fields

Additional features in “old” Web-MARS (2/2)

- **Parameter database**

- **GRIB table based view**
- **Links to IFS documentation**
- **Links to comprehensive list of class, stream and type**

- **Server activity**

- **Show archive activity**
- **Monitor your requests**
- **Learn how the queuing system works**
 - **Reason for queued requests**

Retrieving data - Helpers

Some useful tools

- **grib_ls**, **grib_dump**, ...
- **Metview examiners**
 - `metview4 -e grib [filename]`
 - `metview4 -e bufr [filename]`
- **CDO - Climate Data Operators**

See <https://code.zmaw.de/projects/cdo>

Retrieving data – Conversion to NetCDF

GRIB API tool **grib_to_ncdf**

- To convert a GRIB file to NetCDF format
- GRIB must be a regular lat/lon grid or a regular Gaussian grid
 - i.e. the key "typeOfGrid" should be "regular_ll" or "regular_gg"
- Example
 - > **grib_to_ncdf -o output.nc input.grib1**

See https://software.ecmwf.int/wiki/display/GRIB/grib_to_ncdf

MARS Architecture

- **Client/Server**
- **Protocol: MARS request**
- **Clients, C program + GRIB API + libemos library (Interpolation)**
 - Supercomputers
 - Workstations and Servers
 - Applications like Metview (local / at ECMWF)
 - Remote client for Member States (security mechanism)
 - WebMARS
 - Data Server
 - Web API

MARS Architecture – Servers

- **Reports Database (RDB), on-line observations (for Operations only)**
- **Fields Database (FDB)**
 - Data produced by most recent cycles or experiments
 - Very fast access (on-line data)
 - Suitable for model input
- **ODB database, on-line ODB on supercomputers**
- **Main Archives (6 servers)**
 - Dedicated Linux servers / clustered architecture
 - Terabytes of disk space
 - Tape management SW: HPSS
 - Oracle (Sun) SL8500 Automated Tape Libraries

MARS Architecture - Request execution

- 1) Check syntax (MARS language and request syntax)**
- 2) Print request to be processed**
- 3) Query all Supercomputer's FDB**
- 4) Query main archives (if data not in FDB)**
- 5) Transfer data**
- 6) Post-processing while transferring (if needed)**
- 7) Report on result**

Request execution (1/3)

```
MARS - INFO - **
```

```
MARS - INFO - **
```

```
PPDIR is /ppdir/data/rs60005
```

```
mars - INFO - 20090225.102926 - Welcome to MARS
```

```
retrieve,
```

```
    class      = od,
```

```
    type       = an,
```

```
    expver    = 1,
```

```
    date      = -7,
```

```
    time      = 00/to/18/by/6,
```

```
    param     = t,
```

```
    levtype   = model level,
```

```
    levelist  = 1/to/91,
```

```
    area      = E,
```

```
    grid      = 2.5/2.5,
```

```
    target    = "t.ll"
```

```
mars - INFO - 20090225.102942 - Processing request 1
```

```
mars - WARN - 20090225.102942 - Area not compatible with grid
```

```
mars - WARN - 20090225.102942 - Area changed from 73.5/-27/33/45 to 75/-27.5/32.5/45
```

Request execution (2/3)

```
RETRIEVE,  
  CLASS      = OD,  
  TYPE       = AN,  
  STREAM     = DA,  
  EXPVER    = 0001,  
  REPRES    = SH,  
  LEVTYPE   = ML,  
  LEVELIST   = 1/2/3/4/5/6/7/8/9/10/11/12/13/14/15/16/17/18/19/20/21/22/23/  
24/25/26/27/28/29/30/31/32/33/34/35/36/37/38/39/40/41/42/43/44/45/46/47/48/49/50/5  
1/52/53/54/55/56/57/58/59/60/61/62/63/64/65/66/67/68/69/70/71/72/73/74/75/76/77/78/  
79/80/81/82/83/84/85/86/87/88/89/90/91,  
  PARAM      = 130,  
  DATE       = 20090218,  
  TIME       = 0000/0600/1200/1800,  
  STEP       = 00,  
  DOMAIN    = G,  
  TARGET     = "t.ll",  
  RESOL     = AUTO,  
  AREA       = 75/-27.5/32.5/45,  
  GRID       = 2.5/2.5,  
  PROCESS   = LOCAL
```

Request execution (3/3)

```
mars - INFO - 20090225.102942 - Requesting 364 fields  
819480 FDB; INFO; DB$_ Fields DataBase 4.2  
mars - INFO - 20090225.102942 - Calling mars on 'marsod', callback on 61767  
mars - INFO - 20090225.104347 - Mars client is on ecgate.ecmwf.int (136.156.240.111) 61767  
mars - INFO - 20090225.104347 - Mars server is on hdr16.ecmwf.int (136.156.228.176) 57793  
mars - INFO - 20090225.104347 - Server task is 526 [marsod]  
mars - INFO - 20090225.104347 - Request cost: 364 fields, 445.507 Mbytes online [marsod]  
mars - INFO - 20090225.104347 - Transferring 467148136 bytes  
mars - WARN - 20090225.104348 - INTFB: Resolution automatically set to 63  
mars - INFO - 20090225.104423 - 364 fields retrieved from 'marsod'  
mars - INFO - 20090225.104423 - 364 fields have been interpolated on 'ecgate'  
mars - INFO - 20090225.104423 - Request time: wall: 14 min 42 sec cpu: 12 sec  
mars - INFO - 20090225.104423 - Read from network: 445.51 Mbyte(s) in 24 sec [18.43 Mbyte/sec]  
mars - INFO - 20090225.104423 - Processing in marsod: wall: 14 min 6 sec  
mars - INFO - 20090225.104423 - Visiting marsod: wall: 14 min 42 sec  
mars - INFO - 20090225.104423 - Post-processing: wall: 11 sec cpu: 9 sec  
mars - INFO - 20090225.104423 - Memory used: 13.48 Mbyte(s)  
mars - INFO - 20090225.104423 - No errors reported
```

Retrieving data

Request scheduling

- Queueing system


Priorities: user, request age, request cost (number of tapes and fields)

Data collocation

- MARS tree
- Archive objects (for OD data)

- 1 file per month of AN (1 level type, all times, levels, params)
- 1 file per forecast (1 level type, all steps, levels, params)
- 1 file per EPS (1 level type, all steps, members, levels, params)
- 1 file per month of ERA-40 FC (1 level type, all steps, levels, params)

Retrieving data - MARS tree



Retrieving data - Post-processing

- **Conversions**

- SH → SH (reduced truncation), GG, LL
- GG (reduced) → GG (lower resolution or regular), LL
- LL → LL (lower resolution)

- Sub-area extractions (GG, LL, waves), reduces data volume
- Derived fields (e.g. U and V from vorticity and divergence)
- Rotation

Retrieving data - Post-processing

Truncation before interpolation, reduces necessary resources

Grid increment	Truncation
$2.5 \leq \Delta$	T63
$1.5 \leq \Delta < 2.5$	T106
$0.6 \leq \Delta < 1.5$	T213
$0.4 \leq \Delta < 0.6$	T319
$0.3 \leq \Delta < 0.4$	T511
$0.15 \leq \Delta < 0.3$	T799
$0.09 \leq \Delta < 0.15$	T1279
$0.0 \leq \Delta < 0.09$	T2047

Retrieving data – Efficiency

- **Use local disk (\$SCRATCH)**
- **Estimate amount of data (list command)**
 - Number of fields (up to tens of thousands / request)
 - Data size (up to several Gigabytes / request)
- **Check computer resources: quota, CPU time, ...**
- **Reduce number of tapes involved (better scheduling)**
- **Retrieve as much data from the same tape as possible**
- **Avoid constantly accessing the same tape**
- **Do not create unnecessary sub-archives**

Retrieving data – Data access

- **Archived data**
 - Available to all registered users
- **Current (valid) data, i.e. data for which the value of
 $(\text{DATE} + \text{TIME} + \text{STEP}) + 24 \text{ hours} \geq \text{current date/time}$**
 - Needs special registration
 - Contact your Computing Representative
- **Boundary Conditions Project & COSMO-LEPS**
 - Restricted to participating MS / individual users
- **Restrictions for Observations, TIGGE, EUROSIP...**
- **Data is available according to dissemination schedule**
 - see www.ecmwf.int/services/dissemination/3.1/
- **For time-critical retrievals, use the framework provided**

Data Server – <http://apps.ecmwf.int/datasets/>

- Public distribution of data (licensing depends on datasets)
 - Self-registration
- Based on ecCharts framework
- Datasets

DEMETER

ENSEMBLES

ERA-15

ERA-40, 2.5°

ERA-Interim, full resolution

ERA-20CM (experimental)

MACC reanalysis

YOTC

GEMS

TIGGE

TIGGE-LAM

ISPD v2.2

ICOADS v2.5.1 with interpolated 20CR feedback

Data Server – Web API

- To access ECMWF data servers in batch
- Requirements
 - User account
 - client library, e.g. python
 - API key

See

[https://software.ecmwf.int/wiki/display/WEBAPI/
Accessing+ECMWF+data+servers+in+batch](https://software.ecmwf.int/wiki/display/WEBAPI/Accessing+ECMWF+data+servers+in+batch)

Additional resources

- **MARS documentation**

www.ecmwf.int/publications/manuals/mars/

- **Web-MARS**

apps.ecmwf.int/services/mars/catalogue/

www.ecmwf.int/services/archive/

- **Data Services FAQ**

www.ecmwf.int/products/data/archive/data_faq.html

- **ECMWF forecast products**

www.ecmwf.int/products/forecasts/

- **GRIB API Documentation**

software.ecmwf.int/wiki/display/GRIB/Home

- **IFS Documentation**

www.ecmwf.int/research/ifsdocs/